

Eigenmodes of the Dirac Operator and Chiral Properties of QCD with Sea Quarks

Xiang-Qian Luo ^a

^aDepartment of Physics, Zhongshan University, Guangzhou 510275, China

I describe a mechanism to understand the relation between chiral-symmetry breaking and eigenmodes of the Dirac operator in lattice QCD with Kogut-Susskind sea quarks. It can be shown that if chiral symmetry is spontaneously broken, the eigenvalues λ_i should behave as $z(i)/V$ for large volume V , where $z(i)$ is the i -th zero of the Bessel function. With neither chiral nor λ extrapolation, one can precisely calculate the chiral condensate using only a small set of eigenvalues. Therefore, it is economical and free of systematic uncertainties. I present the first QCD data to support this mechanism and encourage the lattice community to test and use it.

One can not completely understand physics of hadrons without understanding the QCD vacuum, which main properties are confinement and chiral-symmetry breaking.

The chiral order parameter is $\langle\bar{\psi}\psi\rangle$ in the chiral limit. Suppose $\langle\bar{\chi}\chi(m, V)\rangle$ is the one calculated at bare mass m and lattice volume V . Since $\langle\bar{\chi}\chi(0, V)\rangle = 0$, conventionally one has to assume

$$\langle\bar{\psi}\psi\rangle = \lim_{m \rightarrow 0} \lim_{V \rightarrow \infty} \langle\bar{\chi}\chi(m, V)\rangle. \quad (1)$$

Practically, most people do extrapolation only on finite V and very small set of m by means of some unjustified fitting function (linear, quadratic, polynomial or logarithm ...). Unfortunately, such an process is arbitrary and sometimes gives wrong results. Here we provide two known evidences.

(i) QED in 4 dimensions. Non-compact lattice QED experiences second order chiral phase transition at finite some bare coupling constant g , where the chiral condensate vanishes. The critical coupling determined by naive m extrapolation of $\langle\bar{\chi}\chi(m, V)\rangle$ is not necessary correct. For detailed discussions and current status, see [1].

(ii) The one-flavor massless Schwinger model. In the continuum, as is well known, $\langle\bar{\psi}\psi\rangle_{cont} = 0.16e$, where e is the electric charge. As indicated in Ref. [2], the exact result can not be reached at finite bare coupling $g = ae$ by the any extrapolation mentioned above.

In Ref. [3], we proposed a new mechanism to get reliable quantitative information on chiral-

symmetry breaking, based on the computation of the probability distribution function of $\langle\bar{\psi}\psi\rangle$:

$$P(c) = \lim_{N \rightarrow \infty} \langle \delta(c - \frac{1}{N} \sum_x \bar{\psi}(x)\psi(x)) \rangle. \quad (2)$$

Here N is the degrees of freedom, being proportional to V . $P(c)$ is the probability to get the value c for the chiral condensate. If there is exact chiral symmetry in the ground state, $P(c) = \delta(c)$. If chiral symmetry is spontaneously broken, $P(c)$ will be a more complex function.

For Kogut-Susskind fermions, we derive several relations between the chiral condensate and the eigenmodes of the Dirac operator.

(a) The first formula is

$$\langle\bar{\psi}\psi\rangle = \lim_{m \rightarrow 0} \lim_{V \rightarrow \infty} \frac{1}{N} \langle \sum_i^N \frac{2m}{\lambda_i^2 + m^2} \rangle \quad (3)$$

where λ_i is the i -th positive eigenvalue of the massless Dirac matrix. This is the formula many people are currently using. It requires m extrapolation, and may not produce reliable result.

(b) The second formula is

$$\langle\bar{\psi}\psi\rangle = \lim_{\lambda \rightarrow 0} \lim_{V \rightarrow \infty} \frac{\pi}{N} \langle \rho(\lambda) \rangle, \quad (4)$$

where $\rho(\lambda)$ is the eigenmode density. This formula is also known. Since $\rho(0) = 0$ at finite V , and it requires λ extrapolation, which is again arbitrary.

(c) The third formula [3] is

$$\langle \bar{\psi}\psi \rangle = \lim_{V \rightarrow \infty} \sqrt{\frac{4}{N^2} \left\langle \sum_{i=1}^N \frac{1}{\lambda_i^2} \right\rangle}, \quad (5)$$

For its derivation by other methods, see Refs. [4–6]. The advantage over the first two formulae is that no λ extrapolation is necessary. The disadvantage is that all the eigenvalues have to be calculated. When the lattice volume is large, the computational task is huge and not so feasible. Indeed, we have checked that it requires a large lattice volume for getting consistent results.

(d) The fourth formula is

$$\langle \bar{\psi}\psi \rangle = C(i) = \lim_{N \rightarrow \infty} \frac{z(i)}{N} \left\langle \frac{1}{\lambda_i} \right\rangle. \quad (6)$$

Here $z(i)$ is the i -th zero of the Bessel function J_0 . This is new and was originally derived in Ref. [3]. Neither m nor λ extrapolation is necessary. The advantage over the third formula is that only a few smallest eigenvalues are needed for this calculation. The only thing is the finite size analysis which every formula requires to do.

The relation between eigenmodes and chiral-symmetry breaking is clear: if chiral symmetry is spontaneously broken, λ_i should behave as $z(i)/V$ for large V . We have numerically checked [3] the reliability of Eq. (6) in the Schwinger model, where the exact result was reproduced precisely. To our knowledge, this is one of the best results for the lattice Schwinger model!

The most interesting application is QCD. Here I would like to present the first data for QCD with various numbers of quark flavors N_f . In Ref. [7], the configurations were generated and eigenvalues of the fermionic matrix were calculated. With these, it is very easy to evaluate the chiral condensate and compare different methods.

Figure 1 shows the quenched result for $\beta = 5.5088$ and $V = 6^4$ using the first formula: Eq. (3). Only a small set of data can be fitted by a linear function. If one fits only the data for $m \in [0.04, 0.08]$ to the chiral limit, the result is 0.060. But outside this range, either for larger or smaller m , this slope changes and global fitting function can not be easily found. For smaller m ,

the change is very rapid. We attribute it to finite size effects.

Figure 2 shows the quenched result for $\beta = 5.5088$ and $V = 6^4$, but using the fourth formula: Eq. (6). Even on such a small lattice, there is a nice plateau for $i \in [10, 25]$, from which one can reliably get $C(i) = 0.06$. No fitting function is necessary. Of course, detailed finite size analysis remains to be done.

Figures 3, 4, 5, 6 are the results for $N_f = 1, 2, 3, 4$. All these satisfy well Eq. (6).

In conclusion, I have shown a reliable method for investigating the chiral properties and obtaining the chiral condensate from only a small set of λ , without m or λ extrapolation. This method was previously tested in the Schwinger model. In this paper I have presented the first data for QCD. Therefore, I encourage other people in the lattice community to test and use Eq. (6).

This work was supported by the National Natural Science Fund for Distinguished Young Scholars, supplemented by the National Natural Science Foundation, fund for international cooperation and exchange, the Ministry of Education, the Zhongshan University Administrations and Hong Kong Foundation of the Zhongshan University Advanced Research Center. I also thank the Lattice 98 organizers for additional support and assistance. Discussions with V. Azcoiti, T. Chiu, G. DiCarlo, A. Galante, A. Grillo, H. Jirari, H. Kröger, and V. Laliena are appreciated.

REFERENCES

1. A. Kocic, *Nucl. Phys. B(Proc. Suppl.)***34** (1994) 123; V. Azcoiti, *Nucl. Phys. B(Proc. Suppl.)***53** (1997) 148.
2. V. Azcoiti, G. DiCarlo, A. Galante, A. Grillo, V. Laliena, *Phys. Rev. D***50** (1994) 6994.
3. V. Azcoiti, V. Laliena, X.Q. Luo, *Phys. Lett. B***354** (1995) 111.
4. H. Leutwyler, A. Smilga, *Phys. Rev. D***46** (1992) 5607.
5. J. Verbaarschot, *Phys. Lett. B***329** (1994) 351.
6. P. Damgaard, S. Meyer, T. Wettig, contributions to Lattice 98.
7. X.Q. Luo, *Nucl. Phys. B(Proc. Suppl.)***63** (1998) 835.




Figure 1. $\langle \bar{\chi}\chi(m, V) \rangle$ for $N_f = 0$, $\beta = 5.5088$ and $V = 6^4$ using Eq. (3).




Figure 3. $C(i)$ as a function of i for $N_f = 1$, $\beta = 5.3469$ and $V = 6^4$ using Eq. (6).




Figure 2. $C(i)$ as a function of i for $N_f = 0$, $\beta = 5.5088$ and $V = 6^4$ using Eq. (6).




Figure 4. $C(i)$ as a function of i for $N_f = 2$, $\beta = 5.3136$ and $V = 6^4$ using Eq. (6).




Figure 5. $C(i)$ as a function of i for $N_f = 3$, $\beta = 5.2814$ and $V = 6^4$ using Eq. (6).




Figure 6. $C(i)$ as a function of i for $N_f = 4$, $\beta = 5.2500$ and $V = 6^4$ using Eq. (6).